arXiv:cond-mat/0507233v2 [cond-mat.str-el] 10 Apr 2006

Density distributions for trapped one-dimensional spinor gases

Yajiang Hao,[1, 2] Yunbo Zhang,[1] J. Q. Liang,[1] and Shu Chen[2, *]

[1]*Department of Physics and Institute of Theoretical Physics,*
Shanxi University, Taiyuan 030006, P. R. China
[2]*Institute of Physics, Chinese Academy of Sciences, Beijing 100080, P. R. China*

We numerically evaluate the density distribution of a spin-1 bosonic condensate in its ground state within a modified Gross-Pitaevskii theory, which is obtained by the combination of the exact solution of the corresponding integrable model with the local density approximation. Our study reveals that atoms in the $m_F = 0$ state are almost completely suppressed for the anti-ferromagnetic interactions in both weakly and strongly interacting regimes, whereas all three components remain non-vanishing for ferromagnetic interactions. Specially, when the system is in the Tonks-Girardeau (TG) regime, obvious Fermi-like distribution emerges for each component. We also discuss the possible deviation of the spatial distribution from the Fermi-like distribution when the spin-spin interaction is strong enough.

PACS numbers: 03.75.Mn, 03.75.Hh, 67.40.Db

I. INTRODUCTION

The experimental realization of Bose-Einstein condensates (BECs) of trapped alkali atomic clouds [1] has stimulated active studies in many new regimes. From then on, BECs have become a popularly investigated platform for various effects of quantum many-body interaction in strongly correlated systems. When BECs are realized in optical traps rather than in magnetic ones, the spin degrees of freedom in alkali atoms are liberated. As a consequence, such a system with internal degrees of freedom manifests very rich physics and many fascinating phenomena have been observed in spinor condensates, e.g., quantum entanglement of spins, spinor four-wave mixing, and spin domains, etc. [2, 3, 4, 5, 6, 7].

On the other hand, in the research area of low-dimensional physics, the effect of dimensionality reduction in bosonic system has been investigated extensively [8, 9, 10, 11, 12, 13, 14, 15] and is being paid more and more attention [16, 17]. There has been tremendous experimental progress towards the realization of trapped one-dimensional (1D) cold atom systems [18, 19, 20, 21]. Very recently, several groups have reported the observation of a 1D Tonks-Girardeau (TG) gas [22, 23]. An array of 1D quantum gas is obtained by tightly confining the particle motion in two directions to zero point oscillations [24] by means of two-dimensional optical lattice potentials. By loading the condensate in optical lattice and changing the trap intensities, and hence the atomic interaction strength, the atoms can be made to act either like a condensate or like a TG gas. The TG gas provides a textbook example where atom-atom interaction plays a critical role and mean-field theory fails to obtain reasonable results [25]. The dimensionless interaction parameter $\gamma = mg/\hbar^2\rho$ governs the crossover from the weakly interacting condensate to the strongly interacting TG gas, where g is an effective 1D interaction constant, m is the mass of the atom, and ρ is the 1D density of the quantum gas.

Within the mean field theory, Zhang and You recently studied the spin-1 atomic condensate in a cigar-shaped trap by solving an effective quasi-1D nonpolynomial Schrödinger equation [26]. As is well known, the mean field theory fails to work if the system enters the strongly interacting TG regime [27]. It is therefore desirable to investigate the 1D spinor gases within a theoretical approach valid to the strongly interacting TG regime as well as the weakly interacting Thomas-Fermi regime. In this paper, we investigate the ground state properties of the 1D spinor Bose gases under the local density approximation [9, 14, 27] combining with the exact result of the solvable Bose gas model [28, 29, 30]. The dependence of density profile on the magnetization and the effects of many-body interaction are studied in both the weakly interacting regime ($\gamma \ll 1$) and the TG regime ($\gamma \gg 1$). The density profile for the 1D spinor Bose gases in the ground states is then obtained by numerically solving the modified Gross-Pitaevskii Equations (GPEs). Particularly, when the system is in the TG regime, we found that the effect of spin-spin interaction could be dramatic.

The paper is organized as follows. In Sec. II, we give a brief review of the 1D spinor model trapped in an external potential and derive the modified GPEs. In Sec. III, we introduce our numerical procedure for solving the coupled GPEs. Aiming at the realistic systems which may be accessible in experiments, in Sec IV, we present results of the ground state density distributions for the trapped spinor gases in different interacting regimes. A brief summary is given in Sec. V.

II. FORMULATION OF THE MODEL AND METHOD

The s-wave scattering between two identical spin-1 bosons is characterized by the total spin of two collid-

---

[*] Electronic address: schen@aphy.iphy.ac.cn

ing bosons, 0 or 2, and we denote the corresponding scattering lengths by a_0 and a_2. When the cold atoms were trapped intensively in transverse direction with the transverse trapping frequency $\hbar\omega_\perp$ greatly exceeding the chemical potential, the radial motion of atoms is essentially 'frozen out'. The dynamics is thus governed by a 1D Hamiltonian with the effective 1D interaction strength given by [8, 31, 32]

$$\begin{aligned} U_{0,2} &= -\frac{2\hbar^2}{m a_{0,2}^{1D}}, \\ a_{0,2}^{1D} &= -\frac{d_\perp^2}{2a_{0,2}}\left(1-\mathcal{C}\left(a_{0,2}/d_\perp\right)\right), \end{aligned} \tag{1}$$

where $d_\perp = \sqrt{\hbar/m\omega_\perp}$ and $\mathcal{C} \approx 1.4603$. For arbitrary hyperfine states in the hyperfine manifold in which there are two spin-1 atoms, the effective interaction may therefore be written as

$$\begin{aligned} U_{ij} &= \delta\left(x_i - x_j\right)\left(U_0\mathcal{P}_0 + U_2\mathcal{P}_2\right) \\ &= \delta\left(x_i - x_j\right)\left(c_0 + c_2\mathbf{F}_i\cdot\mathbf{F}_j\right), \end{aligned} \tag{2}$$

where the operators $\mathcal{P}_0$ and $\mathcal{P}_2$ project the wave function of a pair of atoms into a state of total spin 0 and 2. The coefficients c_0 and c_2 are related to the effective 1D interaction constants U_0 and U_2 through $c_0 = \frac{U_0+2U_2}{3}$ and $c_2 = \frac{U_2-U_0}{3}$. The ground state of the system is ferromagnetic if $c_2 < 0$, and anti-ferromagnetic if $c_2 > 0$. F_x , F_y and F_z are three spin-1 matrices with the quantization axis taken along the z-axis direction.

For a spin-1 Bose condensate trapped in an external potential $V_{ext}\left(x\right)$ which is independent of the internal states, the Hamiltonian can be expressed in the second quantized form as

$$\begin{aligned} \mathcal{H} &= \int dx \hat{\Psi}_i^+\left(-\frac{\hbar^2}{2m}\frac{d^2}{dx^2} + V_{ext}\left(x\right)\right)\hat{\Psi}_i \\ &+\frac{c_0}{2}\int dx : \left(\hat{\Psi}_i^+\hat{\Psi}_i\right)^2 : \\ &+\frac{c_2}{2}\int dx \hat{\Psi}_k^+\hat{\Psi}_i^+\left(F_\eta\right)_{ij}\left(F_\eta\right)_{kl}\hat{\Psi}_j\hat{\Psi}_l, \end{aligned} \tag{3}$$

where $\hat{\Psi}_i\left(x\right)$ ($\hat{\Psi}_i^\dagger\left(x\right)$) is the field operator that annihilates (creates) an atom in the i-th internal state at location x, and $i = +, 0, -$ denotes the atomic hyperfine state $\left|F = 1, m_F = +1, 0, -1\right\rangle$ respectively. Summation is assumed for repeated indices in the above Hamiltonian and the pair of colons denote the normal-order product. Within the mean field approach the properties of a spinor gas are determined by the following spin-dependent energy functional

$$\begin{aligned} \mathcal{E} &= \int dx\left[\Phi_i^*\left(-\frac{\hbar^2}{2m}\frac{d^2}{dx^2} + V_{ext}\left(x\right)\right)\Phi_i + \rho\epsilon\left(\rho\right)\right] \\ &+\int dx\left[\frac{c_2}{2}\Phi_k^*\Phi_i^*\left(F_\eta\right)_{ij}\left(F_\eta\right)_{kl}\Phi_j\Phi_l\right], \end{aligned} \tag{4}$$

where $\rho = \sum_i \rho_i = \sum_i \left|\Phi_i\right|^2$ and $\epsilon\left(\rho\right) = c_0\rho/2$.

For a one-component Bose gas, it turns out that such a mean field approach works well in the weakly interacting regime but is not able to describe the density distribution correctly in the TG regime. However, a modified Gross-Pitaevskii theory, within which the interaction effect is properly taken into account by using the Lieb-Liniger solution, proves to be able to yield accurate ground-state density distribution even in the Tonks limit [9, 12, 14, 27, 33]. The modified Gross-Pitaevskii theory essentially is based on the local density approximation together with the analytical solution of the homogenous system. Within the local density approximation, the chemical potential of the inhomogeneous gas is determined by the local equilibrium condition,

$$\mu = \mu_{hom}[\rho(x)] + V_{ext}(x),$$

with $\mu_{hom}[\rho]$ being the chemical potential of the corresponding homogeneous system. For the one-component Bose gas, $\mu_{hom}[\rho]$ can be obtained from the solution of the Lieb-Liniger model. In our present problem, the spin exchange strength c_2 is much smaller than the density-density interaction c_0 and the model (3) for $V_{ext}(x) = 0$ and $c_2 = 0$ is the integrable three-component Bose gas model[29, 30]. To make progress, we work in the scheme of modified Gross-Pitaevskii theory [9, 27, 34] and take $\mu_{hom}[\rho]$ from the corresponding integrable model of three-component Bose gas. Effectively, this is equivalent to replacing the energy density $\epsilon\left(\rho\right)$ in Eq. (4) with the energy density of the exactly solvable three-component Bose model, which takes the values in the two limiting cases as following

$$\epsilon\left(\rho\right) = \frac{\hbar^2}{2m}\rho e\left(\gamma\right) = \left\{ \begin{array}{rl} c_0\rho/2, & \gamma \ll 1, \\ \pi^2\hbar^2\rho^2/6m, & \gamma \gg 1. \end{array} \right. \tag{5}$$

The ground state energy density of the three-component Bose model has a similar form as its one-component correspondence, i.e., the Lieb-Liniger model, however, here the density $\rho = \rho_+ + \rho_0 + \rho_-$ is the total density of three components. We note that in Eq. (4) the kinetic energy term is associated with the inhomogeneity of the gas due to external confinement $V_{ext}\left(x\right)$[14, 35] and $\epsilon\left(\rho\right)$ represents essentially the energy density in the homogeneous system, while the last term the spin-spin interaction energy in 1D.

In the weakly interacting regime ($\gamma \ll 1$) the interaction energy will not change the wave function greatly because it is negligibly small compared with the characteristic kinetic energy of an individual atom. In the TG regime ($\gamma \gg 1$), however, the interaction between atoms is so strong that the bosonic atoms behave much like spinless fermions. In both cases, the spin dependent term can be expressed in the explicit form

$$\begin{aligned} &\Phi_k^*\Phi_i^*\left(F_\eta\right)_{ij}\left(F_\eta\right)_{kl}\Phi_j\Phi_l \\ &= \rho_+^2 + \rho_-^2 + 2\rho_0\rho_- + 2\rho_+\rho_0 - 2\rho_+\rho_- \\ &+2\Phi_0^{*2}\Phi_+\Phi_- + 2\Phi_0^2\Phi_-^*\Phi_+^*. \end{aligned} \tag{6}$$

We note that the last two terms in the above equation correspond to the processes that would change the spin states. An atom in the $m_F = 1$ state scatters with another atom in the $m_F = -1$ state, and consequently it produces two atoms in the $m_F = 0$ state or vice versa. Nevertheless, these processes conserve the magnetization of the system $\mathcal{M} = \int dx \langle F \rangle = \int dx \left[\Phi_+^* \Phi_+ - \Phi_-^* \Phi_- \right]$ [26, 36]. In order to obtain the ground state from a global minimization of $\mathcal{E}$ with the constraints on both N and $\mathcal{M}$, we introduce separately Lagrange multiplier B to conserve M and the chemical potential μ to conserve N. The ground state is then determined by a minimization of the free-energy functional $\mathcal{F} = \mathcal{E} - \mu N - B\mathcal{M}$. The dynamics of Φ_i is governed by the coupled GPEs

$$\begin{aligned} i\hbar\partial\Phi_+/\partial t &= \left[H - B + c_2\left(\rho_+ + \rho_0 - \rho_-\right)\right]\Phi_+ + c_2\Phi_0^2\Phi_-^*, \\ i\hbar\partial\Phi_0/\partial t &= \left[H + c_2\left(\rho_+ + \rho_-\right)\right]\Phi_0 + 2c_2\Phi_+\Phi_-\Phi_0^*, \qquad (7) \\ i\hbar\partial\Phi_-/\partial t &= \left[H + B + c_2\left(\rho_- + \rho_0 - \rho_+\right)\right]\Phi_- + c_2\Phi_0^2\Phi_+^*, \end{aligned}$$

with

$$H = -\frac{\hbar^2}{2m}\frac{d^2}{dx^2} + V_{ext}\left(x\right) + \tilde{F}\left(\rho\right) \qquad (8)$$

and

$$\tilde{F}\left(\rho\right) = \frac{\partial}{\partial\rho}\left[\rho\epsilon\left(\rho\right)\right] = \left\{ \begin{array}{cc} c_0\rho, & \gamma \ll 1, \\ \pi^2\hbar^2\rho^2/2m, & \gamma \gg 1. \end{array} \right. \qquad (9)$$

By numerically solving the above equations, we will determine the ground state density distributions for the 1D spinor Bose gases trapped in an harmonic trap $V_{ext}\left(x\right) = \frac{1}{2}m\omega^2x^2$ both in the weakly interacting regime and in the TG regime.

III. NUMERICAL METHOD

To simplify the formalism we choose a slightly different notation. Taking advantage of the fact that all distances and energies in the calculation can be scaled in units of typical length and energy of the external harmonic potential, we introduce the standard length unit

$$a = \sqrt{\frac{\hbar}{m\omega}}, \qquad (10)$$

and rescale the spatial coordinate, the wave function, and the time variable as

$$\begin{aligned} x &= \tilde{x}a, \\ \Phi_i &= \sqrt{\frac{N}{a}}\phi_i, \\ \tilde{\rho}_i &= \left|\phi_i\right|^2, \\ \tilde{\rho} &= \sum_{i=1}^{3}\tilde{\rho}_i, \\ t &= \frac{\tau}{\omega}. \end{aligned}$$



FIG. 1: (Color online) The density profiles of the spin-1 ^{87}Rb condensates in the ground state for the + (solid line), 0 (dashed line) and - (dash-dot lines) components respectively. (a) m=0, Thomas-Fermi regime; (b) m=0.2, Thomas-Fermi regime; (c) m=0, Tonks regime; (d) m=0.2, Tonks regime. In this figure the lengthes are in units of 1.2μm.

The wave function is thus normalized to $\int d\tilde{x} \sum_{i=1}^{3} \left|\phi_i\right|^2 = 1$. With the above changes, our coupled nonlinear Schrödinger equations become

$$\begin{aligned} i\partial\phi_+/\partial\tau &= \left[\tilde{H} - \tilde{B} + g_2\left(\tilde{\rho}_+ + \tilde{\rho}_0 - \tilde{\rho}_-\right)\right]\phi_+ + g_2\phi_0^2\phi_-^*, \\ i\partial\phi_0/\partial\tau &= \left[\tilde{H} + g_2\left(\tilde{\rho}_+ + \tilde{\rho}_-\right)\right]\phi_0 + 2g_2\phi_+\phi_-\phi_0^*, \qquad (11) \\ i\partial\phi_-/\partial\tau &= \left[\tilde{H} + \tilde{B} + g_2\left(\tilde{\rho}_- + \tilde{\rho}_0 - \tilde{\rho}_+\right)\right]\phi_- + g_2\phi_0^2\phi_+^*, \end{aligned}$$

where $\tilde{B} = NB/a\hbar\omega$ and $\tilde{H} = -\left(1/2\right)d^2/d\tilde{x}^2 + \tilde{x}^2/2 + \tilde{F}\left(\tilde{\rho}\right)$ with

$$\tilde{F}\left(\rho\right) = \left\{ \begin{array}{cc} g_0\tilde{\rho} & \gamma \ll 1, \\ \tilde{g}_0\tilde{\rho}^2 & \gamma \gg 1. \end{array} \right.$$

The pair interaction constants are also rescaled as

$$\begin{aligned} g_0 &= Nc_0/a\hbar\omega, \\ \tilde{g}_0 &= N^2\pi^2/2, \\ g_2 &= Nc_2/a\hbar\omega. \end{aligned}$$

By propagating the coupled GPEs Eq. (11) in imaginary time, we obtain the ground state of spin-1 BECs in one dimension. In each propagating step, the wave function ϕ_i is normalized to conserve the atomic number. We ensure the conservation of magnetization $\mathcal{M}$ by adjusting the Lagrange multiplier B. In our procedure the Crank-Nicholson scheme is used to discretize Eq. (11) [37]. We take the initial wave function to be a complex Gaussian with a constant velocity: $exp\left[-x^2/2q - ikx\right]$. Here q and k are adjustable parameters that shall not affect the final converged ground state [36].



FIG. 2: (Color online) The density profiles of the spin-1 ^{23}Na condensates in the ground state for the + (solid line), 0 (dashed line) and - (dash-dot lines) components respectively. (a) m=0, Thomas-Fermi regime; (b) m=0.2, Thomas-Fermi regime; (c) m=0, Tonks regime; (d) m=0.2, Tonks regime. In this figure the lengthes are in units of 7.4μm.



FIG. 3: (Color online) The density profiles of the spin-1 ^{23}Na condensates in the ground state for the + (solid line), 0 (dashed line) and - (dash-dot lines) components respectively. The interaction between atoms is enhanced such that spin interaction parameter g_2 is at the same magnitude as that of the density interaction $\tilde{g}_0$. In this figure the lengthes are in units of 7.4 μm.

IV. DENSITY PROFILES IN THE GROUND STATE

To give a concrete example, we firstly evaluate the density profiles of 1D spinor gases in the ground state for ^{87}Rb (ferromagnetic) with $a_0 = 102a_B$ and $a_2 = 100a_B$ (a_B is the Bohr radius) [38]. By properly choosing the parameters, the system may be prepared either in the weakly interacting regime or in the TG regime. Let us first consider the specific system with typical trap parameters $\omega_x = 0.5$kHz and $\omega_\perp = 50$kHz for $N = 2000$ atoms, in which case the effective interaction strength $\gamma \sim 0.008$ indicating that the system is in the weakly interacting regime. Fig. 1(a) and Fig. 1(b) show the density profiles in units of N/a for m$= \mathcal{M}/N = 0$ and m$= 0.2$, respectively. The density profiles of + component (solid line) and $-$ component (dash-dot lines) superpose on each other exactly as m$= 0$. When the parameters are tuned to $\omega_x = 10$ Hz, $\omega_\perp = 500$kHz and the atomic number to $N = 50$, the effective interaction strength $\gamma \sim 15$ indicates that the system is in the TG regime. The corresponding density profiles are plotted in Fig. 1(c) and Fig. 1(d) for m$= 0$ and m$= 0.2$, respectively. In the TG regime, the atoms distribute uniformly in more extensive area because of the strong interaction in the system. At the boundary the density profiles decrease to zero rapidly, which reminds us the Fermi-Dirac statistical distribution. This imply that the density distribution of the bosonic atoms in the TG regime behave like that of the Fermions.

For the anti-ferromagnetic system we consider a condensate of ^{23}Na with $a_0 = 50a_B$ and $a_2 = 55.1a_B$ [39]. In the weakly interacting regime, the trap frequencies are chosen as $\omega_x = 50$Hz, $\omega_\perp = 10$kHz and $N = 1000$ so that $\gamma \sim 0.001$. The parameters of the system in the TG regime are $\omega_x = 10$Hz, $\omega_\perp = 2000$kHz and $N = 50$ with $\gamma \sim 15$. The density profiles are shown in the Fig. 2 in units of N/a for m$= 0$ and m$= 0.2$, respectively. In the TG regime they exhibit similar fermionization behavior just as in the case of its ferromagnetic counterpart ^{87}Rb. The only difference is that here the population in 0 component remains completely suppressed at zero magnetic field according to the mean field theory. A straight explanation is that atoms prefer to be aligned anti-parallel due to the anti-ferromagnetic spin interaction. On the other hand, atoms in 0 component would collide into pairs with one atom in + component and the other in - component in order to lower the spin interaction energy. This result is in agreement with the calculation for a spinor condensate confined in a spherically symmetric 3D harmonic trap, in which case the condensation occurs for + and - components respectively [40].

For the condensates in the TG regime, the pair interaction coefficient $\tilde{g}_0 = N^2\pi^2/2$ is clearly a constant irrespective of the s -wave scattering length. Should the interaction between atoms be enhanced by the so-called Feshbach resonance, the pair interaction coefficient g_2 could be increased greatly to the magnitude order of the coefficient $\tilde{g}_0$. This enable us to investigate explicitly the effect of the spin-spin interaction. As an example, we consider the system of $N = 15$ ^{23}Na atoms and m$= 0.2$ in the harmonic trap with $\omega_x = 10$Hz and $\omega_\perp = 2000$kHz. This correspond to a spin interaction parameter $g_2 = 500$ while $\tilde{g}_0 = 1110.33$ if the s-wave scattering length a_2 is enhanced to $104.8a_B$. The corresponding density profiles are given in Fig. 3. It is shown that the density profiles are no longer Fermi-like and most atoms are compressed to the narrower range around the center of the harmonic trap. A naive explanation may be that pairs of atoms form singlets due to the strong spin-spin interactions and the effective interaction between the singlets is relatively very weak, therefore the picture of Tonks gas breaks down. Finally, we discuss the conditions under which the above result makes sense. For the ^{23}Na atoms discussed here, the spin exchange energy is always weak since c_2 is two order of magnitude smaller than c_0. Even for the case in which a_2 is enhanced to $104.8a_B$, we have $c_0/c_2 = 4.14$ and the theory presented in this paper still holds. However, for a condensate which enters the regime with g_2 larger than $\tilde{g}_0$, our result is obviously not applicable.

V. SUMMARY

The density profiles of 1D spin-1 Bose gases in the ground state are evaluated in both the weakly interacting regime and the strongly interacting TG regime. The population of atoms in different components depends on the overall magnetization and the (anti-)ferromagnetism of the Bose gases. When the system is in the Tonks regime, the density profiles show obvious Fermi-like distribution. However, for strong enough spin-spin interaction, we observe apparent deviation of the density distribution from the Fermi-like distribution in the TG regime.

Acknowledgments

The authors acknowledge the NSF of China (Grant No. 90203007 and Grant No. 10574150) for financial support. SC is also supported by the "Hundred Talent" program of Chinese Academy of Sciences. We thank L. You, Y. Wang, W.-X. Zhang and W.-D. Li for useful discussions.

---

[1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E.Wieman, E. A. Cornell, Science **269**, 198 (1995); K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, W. Ketterle, Phys. Rev. Lett.**75**, 3969 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett,R. G. Hulet, ibid. **75**, 1687 (1995).

[2] J. O. Andersen, U. Al Khawaja, and H. T. C. Stoof, Phys. Rev. Lett. **88**, 070407 (2002).

[3] T.-L. Ho and S. K. Yip, Phys. Rev. Lett. **84**, 4031 (2000).

[4] T.-L. Ho, Phys. Rev. Lett. **81**, 742 (1998).

[5] C. K. Law, H. Pu, and N. P. Bigelow, Phys. Rev. Lett. **81**, 5257 (1998).

[6] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. 67, 1822 (1998).

[7] W. Zhang, H. Pu, C. Search, and P. Meystre, Phys. Rev. Lett. **88**, 060401 (2002).

[8] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).

[9] V. Dunjko, V. Lorent and M. Olshanii, Phys. Rev. Lett. **86**, 5413 (2001).

[10] D. L. Luxat and A. Griffin, Phys. Rev. A. **67**, 043603 (2003).

[11] P. Pedri, L. Santos, P. Öhberg, and S. Stringari, Phys. Rev. A. **68**, 043601 (2003).

[12] S. Chen and R. Egger, Phys. Rev. A. **68**, 063605 (2003).

[13] K. K. Das, M. D. Girardeau, and E. M.Wright, Phys. Rev. Lett. **89**, 170404 (2002); K. K. Das, G. J. Lapeyre, and E. M. Wright, Phys. Rev. A. **65**, 063603 (2002).

[14] P. Öhberg and L. Santos, Phys. Rev. Lett. **89**, 240402 (2002).

[15] M. D. Girardeau and E. M. Wright, Phys. Rev. Lett. **84**, 5239 (2000); T.-L. Ho and M. Ma, J. Low Temp Phys. **115** , 61 (1999); H. Monien, M. Linn, and N. Elstner, Phys. Rev. A. **58**, R3395 (1998).

[16] A. Recati, P. O. Fedichev, W. Zwerger, and P. Zoller, Phys. Rev. Lett. **90**, 020401 (2003).

[17] G. E. Astrakharchik, D. Blume, S. Giorgini, and B. E. Granger, Phys. Rev. Lett. **92**, 030402 (2004).

[18] A. Görlitz J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle, Phys. Rev. Lett. **87**, 130402 (2001).

[19] H. Moritz, T. Stöferle, M. Köhl, and T. Esslinger, Phys. Rev. Lett. **91**, 250402 (2003); T. Stöferle, H. Moritz, C. Schori, M. Köhl, and T. Esslinger, *ibid.* **92** , 130403 (2004).

[20] S. Richard, F. Gerbier, J. H. Thywissen, M. Hugbart, P. Bouyer, and A. Aspect, Phys. Rev. Lett. **91**, 010405 (2003).

[21] Y.-J. Wang, D. Z. Anderson, V. M. Bright, E. A. Cornell, Q. Diot, T. Kishimoto, M. Prentiss, R. A. Saravanan, S. R. Segal, and S. Wu ., Phys. Rev. Lett. **94**, 090405 (2005).

[22] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G. V. Shlyapnikov, T. W. Hänsch, and I. Bloch, Nature **429**, 277 (2004).

[23] T. Kinoshita, T. Wenger and D. S. Weiss, Science **305**, 1125 (2004).

[24] N. J. van Druten and W. Ketterle, Phys. Rev. Lett. **79**, 549 (1997).

[25] M. D. Girardeau, J. Math. Phys. (N.Y.) **1**, 516 (1960); L. Tonks, Phys. Rev. 50, 955 (1936); V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum inverse scattering method and correlation functions* (Cambridge university press,1993).

[26] W. Zhang and L. You, Phys. Rev. A. **71**, 025603 (2005).

[27] E. B. Kolomeisky, T. J. Newman, J. P. Straley, and X. Qi, Phys. Rev. Lett. **85**, 1146 (2000).

[28] E. H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963); E. H. Lieb, *ibid.* **130**, 1616 (1963).

[29] B. Sutherland, *Beautiful models* (World Scientific, 2004).

[30] Y. Q. Li, S.J. Gu and Z. J. Ying, J. Phys. A: Math. Gen. **36** 2821 (2003).

[31] T. Bergeman, M. G. Moore, and M. Olshanii, Phys. Rev.

Lett. **91**, 163201 (2003); M. D. Girardeau and M. Olshanii, Phys. Rev. A. **70**, 023608 (2002).

[32] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **87**, 050404 (2001); D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, *ibid.* **85**, 3745 (2000).

[33] Y. E. Kim and A. L. Zubarev, Phys. Rev. A. **67**, 015602 (2003).

[34] A. Minguzzi, S. Succi, F. Toschi, M.P. Tosi, P. Vignolo, Phys. Rep. **395**, 223 (2004).

[35] E. H. Lieb, R. Seiringer, and J. Yngvason, Phys. Rev. Lett. **91**, 150401 (2003).

[36] S. Yi, O N . E. Müstecaplioglu, C. P. Sun, and L. You, Phys. Rev. A. **66**, 011601 (2002).

[37] W. H. Press, B. P. Flannery, S. A. Teukolsky, and W. T. Vetterling, *Numerical Recipes in C: The Art of Scientific Computing, 2nd edition* (Cambridge University Press, London, 1992).

[38] E. G. M. van Kempen, S. J. J. M. F. Kokkelmans, D. J. Heinzen, and B. J. Verhaar, Phys. Rev. Lett. **88**, 093201, (2002).

[39] A. Crubellier, O. Dulieu1, F. Masnou-Seeuws, M. Elbs, H. Knöckel, and E. Tiemann, Eur. Phys. J. D **6,** 211 (1999).

[40] W. Zhang, S. Yi and L. You, Phys. Rev. A **70**, 043611 (2004).